
EECS 16A Designing Information Devices and Systems I Discussion 4B
Spring 2021

Recall from lecture the way to compute a determinant of any 2×2 matrix is by using the following formula:

$$\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad \det(\mathbf{A}) = ad - bc$$

1. Mechanical Eigenvalues and Eigenvectors

In each part, find the eigenvalues of the matrix \mathbf{M} and the associated eigenvectors. State if the inverse of \mathbf{M} exists.

(a) $\mathbf{M} = \begin{bmatrix} 0 & 1 \\ -2 & -3 \end{bmatrix}$

Answer:

Let's begin by finding the eigenvalues:

$$\det(\mathbf{A} - \lambda \mathbf{I}) = \det \left(\begin{bmatrix} 0 - \lambda & 1 \\ -2 & -3 - \lambda \end{bmatrix} \right) = 0$$

$$-\lambda(-3 - \lambda) + 2 = 0$$

$$\lambda^2 + 3\lambda + 2 = 0$$

$$(\lambda + 2)(\lambda + 1) = 0$$

$$\lambda = -1, -2$$

$\lambda = -1$:

$$\left[\begin{array}{cc|c} 0 - (-1) & 1 & 0 \\ -2 & -3 - (-1) & 0 \end{array} \right] = \left[\begin{array}{cc|c} 1 & 1 & 0 \\ -2 & -2 & 0 \end{array} \right] \xrightarrow{G.E.} \left[\begin{array}{cc|c} 1 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

$$\begin{array}{l} x_1 + x_2 = 0 \\ x_2 = t \end{array} \implies \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} -t \\ t \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix} t$$

The eigenspace for $\lambda = -1$ is $\text{span} \left\{ \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$.

$\lambda = -2$:

$$\left[\begin{array}{cc|c} 0 - (-2) & 1 & 0 \\ -2 & -3 - (-2) & 0 \end{array} \right] = \left[\begin{array}{cc|c} 2 & 1 & 0 \\ -2 & -1 & 0 \end{array} \right] \xrightarrow{G.E.} \left[\begin{array}{cc|c} 1 & 1/2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

$$\begin{array}{l} x_1 + x_2/2 = 0 \\ x_2 = t \end{array} \implies \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} -t/2 \\ t \end{bmatrix} = \begin{bmatrix} -1/2 \\ 1 \end{bmatrix} t$$

The eigenspace for $\lambda = -2$ is $\text{span} \left\{ \begin{bmatrix} -1/2 \\ 1 \end{bmatrix} \right\}$.

Note that we have no zero eigenvalues, the columns of A are linearly independent, and the determinant of A is non-zero (evaluate our polynomial in λ at $\lambda = 0$). Any of these are equivalent conditions for saying that a square matrix is invertible.

(b) $\mathbf{M} = \begin{bmatrix} -2 & 4 \\ -4 & 8 \end{bmatrix}$

Answer:

Let's begin by finding the eigenvalues:

$$\det(A - \lambda I) = \det \left(\begin{bmatrix} -2 - \lambda & 4 \\ -4 & 8 - \lambda \end{bmatrix} \right) = 0$$

$$(-2 - \lambda)(8 - \lambda) + 16 = 0$$

$$\lambda^2 - 6\lambda = 0$$

$$\lambda(\lambda - 6) = 0$$

$$\lambda = 0, 6$$

$\lambda = 0$:

$$\begin{bmatrix} -2 & 4 & | & 0 \\ -4 & 8 & | & 0 \end{bmatrix} \xrightarrow{G.E.} \begin{bmatrix} 1 & -2 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix}$$

$$\begin{array}{l} x_1 - 2x_2 = 0 \\ x_2 = t \end{array} \implies \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2t \\ t \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix} t$$

The eigenspace for $\lambda = 0$ is $\text{span} \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right\}$.

$\lambda = 6$:

$$\begin{bmatrix} -2 - 6 & 4 & | & 0 \\ 4 & 8 - 6 & | & 0 \end{bmatrix} = \begin{bmatrix} -8 & 4 & | & 0 \\ -4 & 2 & | & 0 \end{bmatrix} \xrightarrow{G.E.} \begin{bmatrix} 1 & -1/2 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix}$$

$$\begin{array}{l} x_1 - x_2/2 = 0 \\ x_2 = t \end{array} \implies \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} t/2 \\ t \end{bmatrix} = \begin{bmatrix} 1/2 \\ 1 \end{bmatrix} t$$

The eigenspace for $\lambda = 6$ is $\text{span} \left\{ \begin{bmatrix} 1/2 \\ 1 \end{bmatrix} \right\}$.

Matrix \mathbf{M} has linearly dependent columns, therefore the inverse \mathbf{M}^{-1} does not exist. Note also that \mathbf{M} has an eigenvalue of 0 so that $N(\mathbf{M})$ contains more than just $\vec{0}$. For this reason also \mathbf{M} is not invertible.

(c) $\mathbf{M} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$

Answer:

Let's begin by finding the eigenvalues:

$$\det(A - \lambda I) = \det \left(\begin{bmatrix} -\lambda & 0 \\ 1 & -\lambda \end{bmatrix} \right) = 0$$

$$\lambda^2 = 0$$

$$\lambda = 0(\times 2)$$

$\lambda = 0$:

$$\begin{bmatrix} 0 & 1 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix} \text{ cannot be further reduced by G.E.}$$

$$x_2 = 0, x_1 = t \implies \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} t \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} t$$

The eigenspace for $\lambda = 0$ is $\text{span}\left\{\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right\}$. Note even though $\lambda = 0$ is an eigenvalue with multiplicity 2 (occurs as a root twice for the characteristic polynomial), the dimension of its eigenspace is only 1. This shows that the number of linearly independent eigenvectors for a given eigenvalue is not necessarily equal to the multiplicity, i.e. the number of times that eigenvalue occurs in the characteristic polynomial.

Matrix \mathbf{M} has a zero column (linearly dependent columns), therefore the inverse \mathbf{M}^{-1} does not exist.

(d) **(PRACTICE)** $\mathbf{M} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$.

Answer: Let's begin by finding the eigenvalues:

$$\det\left(\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix}\right) = \det\left(\begin{bmatrix} -\lambda & -1 \\ 1 & -\lambda \end{bmatrix}\right) = 0$$

$$\lambda^2 + 1 = 0$$

From the above equation, we know that the eigenvalues are $\lambda = i$ and $\lambda = -i$.

For the eigenvalue $\lambda = i$:

$$\begin{aligned} (\mathbf{M} - i\mathbf{I})\vec{x} &= \vec{0} \\ \left(\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} - i\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right)\vec{x} &= \vec{0} \\ \begin{bmatrix} -i & -1 \\ 1 & -i \end{bmatrix}\vec{x} &= \vec{0} \end{aligned}$$

We can also perform Gaussian elimination on matrices with imaginary or complex numbers:

$$\left[\begin{array}{cc|c} -i & -1 & 0 \\ 1 & -i & 0 \end{array}\right] \xrightarrow{G.E.} \left[\begin{array}{cc|c} 1 & -i & 0 \\ 0 & 0 & 0 \end{array}\right] \implies \begin{array}{l} x_1 - ix_2 = 0 \\ x_2 = t \end{array} \implies \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} i \\ 1 \end{bmatrix} t$$

So the eigenspace is $\text{span}\left\{\begin{bmatrix} i \\ 1 \end{bmatrix}\right\}$.

For the eigenvalue $\lambda = -i$:

$$\begin{aligned} (\mathbf{M} + i\mathbf{I})\vec{x} &= \vec{0} \\ \left(\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} + i\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right)\vec{x} &= \vec{0} \\ \left(\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} + \begin{bmatrix} i & 0 \\ 0 & i \end{bmatrix}\right)\vec{x} &= \vec{0} \\ \begin{bmatrix} i & -1 \\ 1 & i \end{bmatrix}\vec{x} &= \vec{0} \end{aligned}$$

$$\left[\begin{array}{cc|c} i & -1 & 0 \\ 1 & i & 0 \end{array}\right] \xrightarrow{G.E.} \left[\begin{array}{cc|c} 1 & i & 0 \\ 0 & 0 & 0 \end{array}\right] \implies \begin{array}{l} x_1 + ix_2 = 0 \\ x_2 = t \end{array} \implies \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} -i \\ 1 \end{bmatrix} t$$

The second eigenspace is $\text{span}\left\{\begin{bmatrix} -i \\ 1 \end{bmatrix}\right\}$.

(e) **(PRACTICE)** $\mathbf{M} = \begin{bmatrix} 1 & 0 \\ 0 & 9 \end{bmatrix}$

Answer:

Let's begin by finding the eigenvalues:

$$\det\left(\begin{bmatrix} 1 & 0 \\ 0 & 9 \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix}\right) = \det\left(\begin{bmatrix} 1-\lambda & 0 \\ 0 & 9-\lambda \end{bmatrix}\right) = 0$$

The determinant of a diagonal matrix is the product of the entries.

$$(1-\lambda)(9-\lambda) = 0$$

From the above equation, we know that the eigenvalues are $\lambda = 1$ and $\lambda = 9$.

For the eigenvalue $\lambda = 1$:

$$\begin{aligned} (\mathbf{M} - 1\mathbf{I})\vec{x} &= \vec{0} \\ \left(\begin{bmatrix} 1 & 0 \\ 0 & 9 \end{bmatrix} - 1\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right)\vec{x} &= \vec{0} \\ \begin{bmatrix} 0 & 0 \\ 0 & 8 \end{bmatrix}\vec{x} &= \vec{0} \end{aligned}$$

From the second equation in the system, $x_2 = 0$, with any solution having the form $\begin{bmatrix} 1 \\ 0 \end{bmatrix} t$ for $t \in \mathbb{R}$. The

eigenspace is thus $\text{span}\left\{\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right\}$.

For the eigenvalue $\lambda = 9$:

$$\begin{aligned} (\mathbf{M} - 9\mathbf{I})\vec{x} &= \vec{0} \\ \left(\begin{bmatrix} 1 & 0 \\ 0 & 9 \end{bmatrix} - 9\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right)\vec{x} &= \vec{0} \\ \left(\begin{bmatrix} 1 & 0 \\ 0 & 9 \end{bmatrix} - \begin{bmatrix} 9 & 0 \\ 0 & 9 \end{bmatrix}\right)\vec{x} &= \vec{0} \\ \begin{bmatrix} -8 & 0 \\ 0 & 0 \end{bmatrix}\vec{x} &= \vec{0} \end{aligned}$$

From the first equation in the system, $x_1 = 0$, so any solution must take the form $\begin{bmatrix} 0 \\ 1 \end{bmatrix} t$ for $t \in \mathbb{R}$. The

eigenspace is $\text{span}\left\{\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right\}$.

The matrix is invertible.

2. Eigenvalues and Special Matrices – Visualization

An eigenvector \vec{v} belonging to a square matrix \mathbf{A} is a nonzero vector that satisfies

$$\mathbf{A}\vec{v} = \lambda\vec{v}$$

where λ is a scalar known as the **eigenvalue** corresponding to eigenvector \vec{v} . Rather than mechanically compute the eigenvalues and eigenvectors, answer each part here by reasoning about the matrix at hand.

- (a) Does the identity matrix in \mathbb{R}^n have any eigenvalues $\lambda \in \mathbb{R}$? What are the corresponding eigenvectors?

Solution/Answer: Multiplying the identity matrix with any vector in \mathbb{R}^n produces the same vector, that is, $\mathbf{I}\vec{x} = \vec{x} = 1 \cdot \vec{x}$. Therefore, $\lambda = 1$. Since \vec{x} can be any vector in \mathbb{R}^n , the corresponding eigenvectors are all vectors in \mathbb{R}^n .

- (b) Does a diagonal matrix $\begin{bmatrix} d_1 & 0 & 0 & \cdots & 0 \\ 0 & d_2 & 0 & \cdots & 0 \\ 0 & 0 & d_3 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & d_n \end{bmatrix}$ in \mathbb{R}^n have any eigenvalues $\lambda \in \mathbb{R}$? What are the corresponding eigenvectors?

Solution/Answer: Since the matrix is diagonal, multiplying the diagonal matrix with any standard basis vector \vec{e}_i produces $d_i\vec{e}_i$, that is, $\mathbf{D}\vec{e}_i = d_i\vec{e}_i$. Therefore, the eigenvalues are the diagonal entries d_i of \mathbf{D} , and the corresponding eigenvector associated with $\lambda = d_i$ is the standard basis vector \vec{e}_i .

- (c) Conceptually, does a rotation matrix in \mathbb{R}^2 by angle θ have any eigenvalues $\lambda \in \mathbb{R}$? For which angles is this the case?

Solution/Answer: In a conceptual sense, there are three cases:

Rotation by 0° : (more accurately, any integer multiple of 360°), which yields a rotation matrix $\mathbf{R} = \mathbf{I}$: This will have one eigenvalue of $+1$ because it doesn't affect any vector ($\mathbf{R}\vec{x} = \vec{x}$). The eigenspace associated with it is \mathbb{R}^2 .

Rotation by 180° : (more accurately, any angle of $180^\circ + n \cdot 360^\circ$ for integer n), which yields a rotation matrix $\mathbf{R} = -\mathbf{I}$: This will have one eigenvalue of -1 because it “flips” any vector ($\mathbf{R}\vec{x} = -\vec{x}$). The eigenspace associated with it is \mathbb{R}^2 .

Any other rotation: there aren't any real eigenvalues. The reason is, if there were any real eigenvalue $\lambda \in \mathbb{R}$ for a non-trivial rotation matrix, it means that we can get $\mathbf{R}\vec{x} = \lambda\vec{x}$ for some $\vec{x} \neq \vec{0}$, which means that by rotating a vector, we scaled it. This is a contradiction (again, unless $\mathbf{R} = \mathbf{I}$). Refer to Figure 1 for a visualization.




Figure 1: Rotation will never scale any non-zero vector (by a real number) unless it is rotation by an integer multiple of 360° (identity matrix) or the rotation angle is $\theta = 180^\circ + n \cdot 360^\circ$ for any integer n ($-\mathbf{I}$).

- (d) **(PRACTICE)** Now let us mechanically compute the eigenvalues of the rotation matrix in \mathbb{R}^2 . Does it agree with our findings above? As a refresher, the rotation matrix \mathbf{R} has the following form:

$$\mathbf{R} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

Solution/Answer: Using our known determinant formula for 2×2 matrices $\det(A) = ad - bc$ we can compute the characteristic polynomial

$$\det(\mathbf{R} - \lambda \mathbf{I}) = \det \begin{bmatrix} \cos(\theta) - \lambda & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) - \lambda \end{bmatrix} = \cos(\theta)^2 + \sin(\theta)^2 - 2\cos(\theta)\lambda + \lambda^2 \equiv 0$$

From here we can first simplify $1 = \cos(\theta)^2 + \sin(\theta)^2$ and then use the quadratic formula to attain the two possible λ values.

$$\lambda = \cos(\theta) \pm \sqrt{\cos(\theta)^2 - 1} = \cos(\theta) \pm i\sqrt{1 - \cos(\theta)^2} = \cos(\theta) \pm i\sqrt{\sin(\theta)^2}$$

In exponential phase notation we can write the two eigenvalues more concisely: $\lambda = e^{\pm i\theta}$

- (e) Does the reflection matrix \mathbf{T} across the x-axis in $\mathbb{R}^{2 \times 2}$ have any eigenvalues $\lambda \in \mathbb{R}$?

$$\mathbf{T} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

Solution/Answer: Yes, both $+1$ and -1 . Mechanically, we could go through the methods we have learned for attaining a characteristic polynomial from $\det(T - \lambda I) = (1 - \lambda)(-1 - \lambda) - (0)(0)$ and recalling our eigenvalues are the roots of this polynomial (the values where this polynomial is zero). This works because matrix $T - \lambda I$ only has a nonempty null space when its determinant is zero!

$$\det(T - \lambda I) = \lambda^2 - 1 \equiv 0 \rightarrow \lambda = \pm 1$$

Conceptually, we can reason that a vector along the x-axis will be unaffected by \mathbf{T} (in this case $\lambda = +1$), whereas a vector along the y-axis gets perfectly flipped by \mathbf{T} (in this case $\lambda = -1$)

NOTE: A 2×2 reflection matrix always has $\lambda = \pm 1$, REGARDLESS of the axis of reflection. Why? Reflecting any vector that is on the reflection axis will not affect it (eigenvalue $+1$). Reflecting any

vector orthogonal (perpendicular) to the reflection axis will just “flip it/negate it” (eigenvalue -1). In other words, the set of vectors that lie along the axis of reflection is the eigenspace associated with the eigenvalue $+1$ and the set of vectors orthogonal to the axis of reflection is the eigenspace associated with the eigenvalue -1 .

- (f) If a matrix \mathbf{M} has an eigenvalue $\lambda = 0$, what does this say about its null space? What does this say about the solutions of the system of linear equations $\mathbf{M}\vec{x} = \vec{b}$?

Solution/Answer: $N(A)$ is not just $\vec{0}$ as we have some $\vec{v} \neq \vec{0}$ satisfying $A\vec{v} = \lambda\vec{v}$. Another way we can state this is that $\dim(N(A)) > 0$.

Thus we can imagine if $\mathbf{M}\vec{x} = \vec{b}$ has a solution then $\mathbf{M}(\vec{x} + \vec{v}) = \vec{b}$ also solves the system, hence there are infinite solutions. Yet we also know that a nonzero null space means \mathbf{M} has linearly dependent columns, so the vector \vec{b} could lie outside of this span in which case there is no solution.

In summary, there are either infinite or no solutions to the system of equations $\mathbf{M}\vec{x} = \vec{b}$

- (g) **(Practice)** Does the matrix $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ have any eigenvalues $\lambda \in \mathbb{R}$? What are the corresponding eigenvectors?

Solution/Answer:

Note that the matrix has linearly dependent columns. Therefore, according to part (f), one eigenvalue is $\lambda = 0$. The corresponding eigenvector, which is equivalent to the basis vector for the null space, is

$\begin{bmatrix} 1 \\ -1 \end{bmatrix}$. The other eigenvalue is, by inspection, $\lambda = 1$ with the corresponding eigenvector $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$ because

$$\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}.$$

3. Steady and Unsteady States

(a) You're given the matrix \mathbf{M} :

$$\mathbf{M} = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & -\frac{1}{2} \\ 0 & 1 & -2 \\ 0 & 0 & 2 \end{bmatrix}$$

Which generates the next state of a physical system from its previous state: $\vec{x}[k+1] = \mathbf{M}\vec{x}[k]$. (\vec{x} could describe either people or water.) Find the eigenspaces associated with the following eigenvalues:

- i. $\text{span}(\vec{v}_1)$, associated with $\lambda_1 = 1$
- ii. $\text{span}(\vec{v}_2)$, associated with $\lambda_2 = 2$
- iii. $\text{span}(\vec{v}_3)$, associated with $\lambda_3 = \frac{1}{2}$

Solution/Answer:

i. $\lambda = 1$:

$$\left[\mathbf{M} - \mathbf{I} \mid \vec{0} \right] = \left[\begin{array}{ccc|c} -\frac{1}{2} & \frac{1}{2} & -\frac{1}{2} & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right] \xrightarrow{G.E.} \left[\begin{array}{ccc|c} 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$\vec{v}_1 = \alpha \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \alpha \in \mathbb{R}$$

ii. $\lambda = 2$

$$\left[\mathbf{M} - 2\mathbf{I} \mid \vec{0} \right] = \left[\begin{array}{ccc|c} -\frac{3}{2} & \frac{1}{2} & -\frac{1}{2} & 0 \\ 0 & -1 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \xrightarrow{G.E.} \left[\begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$\vec{v}_2 = \beta \begin{bmatrix} -1 \\ -2 \\ 1 \end{bmatrix}, \beta \in \mathbb{R}$$

iii. $\lambda = \frac{1}{2}$

$$\left[\mathbf{M} - \frac{1}{2}\mathbf{I} \mid \vec{0} \right] = \left[\begin{array}{ccc|c} 0 & \frac{1}{2} & -\frac{1}{2} & 0 \\ 0 & \frac{1}{2} & -2 & 0 \\ 0 & 0 & \frac{3}{2} & 0 \end{array} \right] \xrightarrow{G.E.} \left[\begin{array}{ccc|c} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$\vec{v}_3 = \gamma \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \gamma \in \mathbb{R}$$

Solution/Answer: (8 min)

(b) Define $\vec{x} = \alpha\vec{v}_1 + \beta\vec{v}_2 + \gamma\vec{v}_3$, a linear combination of the eigenvectors. For each of the cases in the table, determine if

$$\lim_{n \rightarrow \infty} \mathbf{M}^n \vec{x}$$

converges. If it does, what does it converge to?

α	β	γ	Converges?	$\lim_{n \rightarrow \infty} \mathbf{M}^n \vec{x}$
0	0	$\neq 0$		
0	$\neq 0$	0		
0	$\neq 0$	$\neq 0$		
$\neq 0$	0	0		
$\neq 0$	0	$\neq 0$		
$\neq 0$	$\neq 0$	0		
$\neq 0$	$\neq 0$	$\neq 0$		

Answer:

$$\begin{aligned}
 \mathbf{M}^n \vec{x} &= \mathbf{M}^n (\alpha \vec{v}_1 + \beta \vec{v}_2 + \gamma \vec{v}_3) \\
 &= \alpha \mathbf{M}^n \vec{v}_1 + \beta \mathbf{M}^n \vec{v}_2 + \gamma \mathbf{M}^n \vec{v}_3 \\
 &= 1^n \alpha \vec{v}_1 + 2^n \beta \vec{v}_2 + \left(\frac{1}{2}\right)^n \gamma \vec{v}_3
 \end{aligned}$$

α	β	γ	Converges?	$\lim_{n \rightarrow \infty} \mathbf{M}^n \vec{x}$
0	0	$\neq 0$	Yes	$\vec{0}$
0	$\neq 0$	0	No	-
0	$\neq 0$	$\neq 0$	No	-
$\neq 0$	0	0	Yes	$\alpha \vec{v}_1$
$\neq 0$	0	$\neq 0$	Yes	$\alpha \vec{v}_1$
$\neq 0$	$\neq 0$	0	No	-
$\neq 0$	$\neq 0$	$\neq 0$	No	-