



## 2. Exploring Column Spaces and Null Spaces

- The **column space** is the **span** of the column vectors of the matrix.
- The **null space** is the set of input vectors that when multiplied with the matrix result in the zero vector.

For the following matrices, answer the following questions:

- What is the column space of  $\mathbf{A}$ ? What is its dimension?
- What is the null space of  $\mathbf{A}$ ? What is its dimension?
- Are the column spaces of the row reduced matrix  $\mathbf{A}$  and the original matrix  $\mathbf{A}$  the same?
- Do the columns of  $\mathbf{A}$  span  $\mathbb{R}^2$ ? Do they form a basis for  $\mathbb{R}^2$ ? Why or why not?

(a)  $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$

**Answer:** Column space:  $\text{span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$

Null space:  $\text{span} \left\{ \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$

The matrix is already row reduced. The column spaces of the row reduced matrix and the original matrix are the same.

The column space does not span  $\mathbb{R}^2$  and thus are not a basis for  $\mathbb{R}^2$ .

(b)  $\begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}$

**Answer:**

Column space:  $\text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$

Null space:  $\text{span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$

The two column spaces are not the same.

Not a basis for  $\mathbb{R}^2$ .

(c)  $\begin{bmatrix} 1 & 2 \\ -1 & 1 \end{bmatrix}$

**Answer:**

Column space:  $\mathbb{R}^2$

Null space:  $\text{span} \left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$

The two column spaces are the same as the column span  $\mathbb{R}^2$ , since they are two independent vectors. This is a basis for  $\mathbb{R}^2$ .

(d)  $\begin{bmatrix} -2 & 4 \\ 3 & -6 \end{bmatrix}$

**Answer:**

Column space:  $\text{span} \left\{ \begin{bmatrix} 1 \\ -\frac{3}{2} \end{bmatrix} \right\}$

Null space:  $\text{span} \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right\}$

The two column spaces are not the same. We can also see that one of the columns is a scaled version of the other, and therefore, they are linearly dependent vectors.

Not a basis for  $\mathbb{R}^2$ .

(e)  $\begin{bmatrix} 1 & -1 & -2 & -4 \\ 1 & 1 & 3 & -3 \end{bmatrix}$

**Answer:**

- i. The column space of the columns is  $\mathbb{R}^2$ . The columns of  $\mathbf{A}$  do not form a basis for  $\mathbb{R}^2$ . This is because the columns of  $\mathbf{A}$  are linearly dependent.
- ii. The following algorithm can be used to solve for the null space of a matrix. The procedure is essentially solving the matrix-vector equation  $\mathbf{A}\vec{x} = \vec{0}$  by performing Gaussian elimination on  $\mathbf{A}$ . We start by performing Gaussian elimination on matrix  $\mathbf{A}$  to get the matrix into upper-triangular form.

$$\begin{aligned} \begin{bmatrix} 1 & -1 & -2 & -4 \\ 1 & 1 & 3 & -3 \end{bmatrix} &\sim \begin{bmatrix} 1 & -1 & -2 & -4 \\ 0 & 2 & 5 & 1 \end{bmatrix} \\ &\sim \begin{bmatrix} 1 & -1 & -2 & -4 \\ 0 & 1 & \frac{5}{2} & \frac{1}{2} \end{bmatrix} \\ &\sim \begin{bmatrix} 1 & 0 & \frac{1}{2} & -\frac{7}{2} \\ 0 & 1 & \frac{5}{2} & \frac{1}{2} \end{bmatrix} \text{ reduced row echelon form} \end{aligned}$$

$$x_1 + \frac{1}{2}x_3 - \frac{7}{2}x_4 = 0$$

$$x_2 + \frac{5}{2}x_3 + \frac{1}{2}x_4 = 0$$

$x_3$  is free and  $x_4$  is free

Now let  $x_3 = s$  and  $x_4 = t$ . Then we have:

$$\begin{aligned}x_1 + \frac{1}{2}s - \frac{7}{2}t &= 0 \\x_2 + \frac{5}{2}s + \frac{1}{2}t &= 0\end{aligned}$$

Now writing all the unknowns  $(x_1, x_2, x_3, x_4)$  in terms of the dummy variables:

$$\begin{aligned}x_1 &= -\frac{1}{2}s + \frac{7}{2}t \\x_2 &= -\frac{5}{2}s - \frac{1}{2}t \\x_3 &= s \\x_4 &= t\end{aligned}$$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} -\frac{1}{2}s + \frac{7}{2}t \\ -\frac{5}{2}s - \frac{1}{2}t \\ s \\ t \end{bmatrix} = \begin{bmatrix} -\frac{1}{2}s \\ -\frac{5}{2}s \\ s \\ 0 \end{bmatrix} + \begin{bmatrix} \frac{7}{2}t \\ -\frac{1}{2}t \\ 0 \\ t \end{bmatrix} = s \begin{bmatrix} -\frac{1}{2} \\ -\frac{5}{2} \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} \frac{7}{2} \\ -\frac{1}{2} \\ 0 \\ 1 \end{bmatrix}$$

So every vector in the null space of  $\mathbf{A}$  can be written as follows:

$$\text{Null space}(\mathbf{A}) = s \begin{bmatrix} -\frac{1}{2} \\ -\frac{5}{2} \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} \frac{7}{2} \\ -\frac{1}{2} \\ 0 \\ 1 \end{bmatrix}$$

Therefore the null space of  $\mathbf{A}$  is

$$\text{span} \left\{ \left[ \begin{bmatrix} -\frac{1}{2} \\ -\frac{5}{2} \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} \frac{7}{2} \\ -\frac{1}{2} \\ 0 \\ 1 \end{bmatrix} \right] \right\}$$

$\mathbf{A}$  has a 2-dimensional null space.

- iii. In this case, the column space of the row reduced matrix is also  $\mathbb{R}^2$ , but this need not be true in general.
  - iv. No, the columns of  $\mathbf{A}$  do not form a basis for  $\mathbb{R}^2$ .
- (f) What do you notice about the relationship between the dimension of the column space, the dimension of the null space, and their sum in all of these matrices?

**Answer:**

The dimension of the column space indicates the number of linearly independent columns in the matrix, while the dimension of the null space indicates the number of linearly dependent columns in the matrix. In each of the above matrices, notice that summing the dimension of the column space and null space results in the total number of columns of the original matrix. This concept forms the rank-nullity theorem.